

SPECIALTY PRODUCTS: ACTIVATED PROTHROMBIN COMPLEXES, BY-PASSING AGENTS FOR TREATMENT OF: FACTOR VIII or IX INHIBITORS; ACQUIRED HEMOPHILIA; ALTERED HEMOSTASIS DUE TO LIVER OR MULTIPLE ORGAN DYSFUNCTIONS

	FEIBA-VH	NOVOSEVEN RT
MANUFACTURER	BAXTER	NOVO NORDISK
CATEGORY	ACTIVATED PCC (prothrombin concentrate complex)	"By passing activity"
CONTENTS	Partial activation during purification process, results in "inhibitor bypassing activity"	Activated VIIa
FACTOR II	yes	
FACTOR VII	activated form	
FACTOR IX	yes	
FACTOR X	yes	
SOURCE MATERIAL	Pooled human plasma	Recombinant, baby hamster kidney cell lines (BHK cells)
DOSING	75-100 units/kg, repeated twice daily if necessary	90 micrograms / kg repeated every 2 hours until bleeding stops
	DO NOT exceed 200 units/kg/day	(Dosage range = 30-120 micrograms/kg) Most hemorrhages stop at 3-4 infusions
VIRAL ATTENUATION METHODS	vapor heat	none specifically stated, but uses chromatographic processes
BOX CONTENTS	Diluent: Sterile Water 20 ml (high); 50ml (ultra high); Baxject mix device	Diluent: buffered Sterile Water; 1mg vial = 1.1ml; 2mg vial = 2.1ml; 5mg vial = 5.1ml; all yield concentration of 1mg/ml
		Infusion Kit (1 per dose): 5 vial adapters, 10cc syringe, 30cc syringe, alcohol swabs, 23g winged infusion set
STORAGE REQUIREMENTS	Refrigerate	Room Temperature
COMMENTS	Administer at rate of 2units/kg per minute.	To avoid product waste, dose should be calculated to nearest complete vial size.
		Recombinant technology may be ONLY product of choice for patients of Jehovah Witness faith.